









APPLICATION RATES BY CROPS

	CROPS	DOSE L/ha	APPLICATION PERIOD
	Alfalfa	3 L/ ha	10 days after each cut.
Q	Cotton	3 L/ ha	1st. Application at the beginning of the delimitation of the plot segment. 2nd . 15 days later. 3rd . 15 days after the 2nd.
১ ৯	Rice, Barley, Corn, Oats	3 L/ ha	1st. Application during cane development. 2nd. Application when the plant sheaths have thickened. 3rd. At flowering.
*	Chili peppers	3 L/ ha	1st. Application at flowering. 2nd. 15 days after. 3rd. 15 days after the 2nd.
***	Beans	3 L/ ha	1st. Application at flowering. 2nd. 15 days after
*	Apples	3 L/ ha	1st. Application after pruning 2nd. 20 days after 3rd. 20 days after the 2nd.
	Watermelons	2.5 L/ ha	1st. Application at flowering. 2nd. 15 days after 3rd. 10 days after the 2nd.
**	Soybeans	3 L/ ha	1st. Application at flowering. 2nd. 15 days after 3rd. 10 days after the 2nd.
	Melons	3 L/ ha	1st. Application at flowering. 2nd. 15 days after 3rd. 10 days after the 2nd.
(3)	Cucumbers	3 L/ ha	1st. Application at flowering. 2nd. 15 days after 3rd. 10 days after the 2nd.
	Wheat	2.5 L/ ha	1st. Application when the plant sheaths have thickened. 2nd. Application at flowering.
*	Grapes	3 L/ ha	1st. Application at flowering. 2nd. 15 days after 3rd. 15 days after the 2nd.
	Tomatoes	3 L/ ha	1st. Application at flowering. Repeat every two weeks.











