



Provides the calcium and boron necessary for the proper development of crops.

Feuillaje® CABO provides and corrects calcium and boron deficiencies in a synergic and effective way to improve flowers and fruits thanks to its source and concentration of elements, as well as its fast assimilation technology.

Benefits:

- Reduces fruit softening and deterioration.
- Greater consistency and quality of fruits and tissues.
- Increases fertilization.
- Decreases the incidence of post - harvest diseases.





Plentiful and turgid structures.



APPLICATION RATES BY CROPS

CROP	DOSE L / ha	APPLICATION PERIOD
 Corn, sorghum, wheat, barley, oats	1.5 L/ha	Apply at the 4-leaf stage and repeat during grain filling.
 Beans, chickpeas, fava beans, peas	1-2 L/ha	Apply at the beginning of flowering and repeat every 15 days.
 Tomato, green tomato and chili	2 - 5 L/ha	Apply at the 4-leaf stage and repeat during grain filling.
 Melon, watermelon, pumpkin and cucumber	2 - 4 L/ha	During flowering, repeat at fruit set and fruit development.
 Avocado	3 - 6 L/ha	At full flowering, repeat every 15 - 20 days, making 5 - 6 applications throughout the cycle.
 Celery, lettuce, cabbage and cauliflower	2 - 4 L/ha	Apply every 10 days, starting mid-cycle and continuing until harvest.
 Potato	2 - 4 L/ha	Apply at the onset of tuberization. Repeat after 15 and 30 days.
 Alfalfa	1 - 2 L/ha	Apply when the plant has 4 to 6 true leaves and after each cutting.
 Sugar cane	1.5 L/ha	Apply during vegetative growth.

Feuillaje® CABO stimulates the plant's natural defense compounds, increasing tolerance to adverse abiotic factors.



agroenzymas.com



comunicacion@retenum.com

